

1 What is claimed is:

- 2
- 3 1. An isolated nucleic acid molecule selected from the group consisting of:
- 4 a) a nucleic acid molecule comprising a nucleotide sequence of SEQ ID NO:1,
- 5 or SEQ ID NO:3;
- 6 b) a nucleic acid molecule which encodes a polypeptide comprising the amino
- 7 acid sequence of SEQ ID NO:2;
- 8 c) a nucleic acid molecule which encodes a fragment of a polypeptide
- 9 comprising the amino acid sequence of SEQ ID NO:2, wherein the fragment comprises at
- 10 least 120 contiguous amino acids of SEQ ID NO: 2; and
- 11 d) a nucleic acid molecule which encodes a naturally occurring allelic variant of
- 12 a polypeptide comprising the amino acid sequence of SEQ ID NO:2, wherein the nucleic
- 13 acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO: 1, 3, or a
- 14 complement thereof, under stringent conditions.
- 15
- 16 2. The isolated nucleic acid molecule of claim 1, which is selected from the
- 17 group consisting of:
- 18 a) a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, SEQ ID
- 19 NO:3; and
- 20 b) a nucleic acid molecule which encodes a polypeptide comprising the amino
- 21 acid sequence of SEQ ID NO:2.
- 22
- 23 3. The nucleic acid molecule of claim 1 further comprising vector nucleic acid
- 24 sequences.
- 25
- 26 4. The nucleic acid molecule of claim 1 further comprising nucleic acid
- 27 sequences encoding a heterologous polypeptide.
- 28
- 29 5. A host cell which contains the nucleic acid molecule of claim 1.
- 30
- 31 6. The host cell of claim 5 which is a mammalian host cell.
- 32

1 7. A non-human mammalian host cell containing the nucleic acid molecule of
2 claim 1.

3
4 8. An isolated polypeptide selected from the group consisting of:

5 a) a polypeptide which is encoded by a nucleic acid molecule comprising a
6 nucleotide sequence which is at least 95% identical to a nucleic acid comprising the
7 nucleotide sequence of SEQ ID NO: 1, SEQ ID NO:3, or a complement thereof.

8 b) a naturally occurring allelic variant of a polypeptide comprising the amino
9 acid sequence of SEQ ID NO:2, wherein the polypeptide is encoded by a nucleic acid
10 molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO: 1, SEQ ID
11 NO:3, or a complement thereof under stringent conditions; and

12 c) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID
13 NO:2, wherein the fragment comprises at least 120 contiguous amino acids of SEQ ID
14 NO:2.

15
16 9. The isolated polypeptide of claim 8 comprising the amino acid sequence of
17 SEQ ID NO:2.

18
19 10. The polypeptide of claim 8 further comprising heterologous amino acid
20 sequences.

21
22 11. An antibody which selectively binds to a polypeptide of claim 8.

23
24 12. A method for producing a polypeptide selected from the group consisting of:

25 a) a polypeptide comprising the amino acid sequence of SEQ ID NO:2;

26 b) a polypeptide comprising a fragment of the amino acid sequence of SEQ ID
27 NO:2, wherein the fragment comprises at least 120 contiguous amino acids of SEQ ID
28 NO:2; and

29 c) a naturally occurring allelic variant of a polypeptide comprising the amino
30 acid sequence of SEQ ID NO:2, wherein the polypeptide is encoded by a nucleic acid
31 molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID
32 NO:3, or a complement thereof under stringent conditions;

1 comprising culturing the host cell of claim 5 under conditions in which the nucleic
2 acid molecule is expressed.

3
4 13. A method for detecting the presence of a polypeptide of claim 8 in a sample,
5 comprising:

6 a) contacting the sample with a compound which selectively binds to a
7 polypeptide of claim 8; and

8 b) determining whether the compound binds to the polypeptide in the sample.
9

10 14. The method of claim 13, wherein the compound which binds to the
11 polypeptide is an antibody.
12

13 15. A kit comprising a compound which selectively binds to a polypeptide of
14 claim 8 and instructions for use.
15

16 16. A method for detecting the presence of a nucleic acid molecule of claim 1 in
17 a sample, comprising the steps of:

18 a) contacting the sample with a nucleic acid probe or primer which selectively
19 hybridizes to the nucleic acid molecule; and

20 b) determining whether the nucleic acid probe or primer binds to a nucleic acid
21 molecule in the sample.
22

23 17. The method of claim 16, wherein the sample comprises mRNA molecules
24 and is contacted with a nucleic acid probe.
25

26 18. A kit comprising a compound which selectively hybridizes to a nucleic acid
27 molecule of claim 1 and instructions for use.
28

29 19. A method for identifying a compound which binds to a polypeptide of claim
30 8 comprising the steps of:

31 a) contacting a polypeptide, or a cell expressing a polypeptide of claim 8 with a
32 test compound; and

33 b) determining whether the polypeptide binds to the test compound.

1
2 20. The method of claim 19, wherein the binding of the test compound to the
3 polypeptide is detected by a method selected from the group consisting of:
4 a) detection of binding by direct detecting of test compound/polypeptide
5 binding;
6 b) detection of binding using a competition binding assay;
7 c) detection of binding using an assay for 25466-mediated signal transduction.

8
9 21. A method for modulating the activity of a polypeptide of claim 8 comprising
10 contacting a polypeptide or a cell expressing a polypeptide of claim 8 with a compound
11 which binds to the polypeptide in a sufficient concentration to modulate the activity of the
12 polypeptide.

13
14 22. A method for identifying a compound which modulates the activity of a
15 polypeptide of claim 8, comprising:
16 a) contacting a polypeptide of claim 8 with a test compound; and
17 b) determining the effect of the test compound on the activity of the polypeptide
18 to thereby identify a compound which modulates the activity of the polypeptide.